

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
25 November 2004 (25.11.2004)

PCT

(10) International Publication Number
WO 2004/101617 A1

(51) International Patent Classification⁷: **C07K 14/715**,
14/54, 14/47, A61K 38/20

(21) International Application Number:
PCT/EP2004/005073

(22) International Filing Date: 11 May 2004 (11.05.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
03101326.1 13 May 2003 (13.05.2003) EP

(71) Applicant (for all designated States except US): **APPLIED RESEARCH SYSTEMS ARS HOLDING N.V.**
[NL/NL]; Pietermaai 15, NL- Curacao (NL).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **ALTAROCCA, Valter** [IT/IT]; Via P.G. Frassati 19 sc.B Santa Mariadelle Mole, Marino, I-00047 Rome (IT). **PEZZOTTI, Anna, R.** [IT/IT]; Via Gubbio, 42, I-00181 Rome (IT).

(74) Agent: **SERONO INTERNATIONAL S.A. INTELLECTUAL PROPERTY**; 12, chemin des Aulx, CH-1228 Plan-Les-Quates (CH).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: ACTIVE VARIANTS OF THE IL-18 BINDING PROTEIN AND MEDICAL USES THEREOF

(57) Abstract: The invention relates to active fragments of the IL-18 binding protein, to pharmaceutical compositions comprising such active fragments, and to medical uses thereof.



ACTIVE VARIANTS OF THE IL-18 BINDING PROTEIN AND MEDICAL USES
THEREOF

FIELD OF THE INVENTION

5 The present invention relates to new interleukin 18 binding proteins (IL-18BPs). The invention further relates to pharmaceutical compositions comprising such IL-18BPs, to nucleic acids encoding such IL-18BPs and to medical uses of said IL-18BPs.

BACKGROUND OF THE INVENTION

10 In 1989, an endotoxin-induced serum activity that induced interferon- γ (IFN- γ) obtained from mouse spleen cells was described (Nakamura et al., 1989). This serum activity functioned not as a direct inducer of IFN- γ but rather as a co-stimulant together with IL-2 or mitogens. An attempt to purify the activity from post-endotoxin mouse serum revealed an apparently homogeneous 50-55 kDa protein. Since other cytokines
15 can act as co-stimulants for IFN- γ production, the failure of neutralizing antibodies to IL-1, IL-4, IL-5, IL-6, or TNF to neutralize the serum activity suggested it was a distinct factor. In 1995, the same scientists demonstrated that the endotoxin-induced co-stimulant for IFN- γ production was present in extracts of livers from mice preconditioned with *P. acnes* (Okamura et al., 1995). In this model, the hepatic
20 macrophage population (Kupffer cells) expand and in these mice, a low dose of bacterial lipopolysaccharide (LPS), which in non-preconditioned mice is not lethal, becomes lethal. The factor, named IFN- γ -inducing factor (IGIF) and later designated interleukin-18 (IL-18), was purified to homogeneity from 1,200 grams of *P. acnes*-
25 treated mouse livers. Degenerate oligonucleotides derived from amino acid sequences of purified IL-18 were used to clone a murine IL-18 cDNA. IL-18 is an 18-19 kDa protein of 157 amino acids, which has no obvious similarities to any peptide in the databases. Messenger RNAs for IL-18 and interleukin-12 (IL-12) are readily detected in Kupffer cells and activated macrophages. Recombinant IL-18 induces IFN-gamma more
30 potently than does IL-12, apparently through a separate pathway (Micallef et al., 1996). Similar to the endotoxin-induced serum activity, IL-18 does not induce IFN- γ by itself, but functions primarily as a co-stimulant with mitogens or IL-2. IL-18 enhances T cell proliferation, apparently through an IL-2-dependent pathway, and enhances Th1

cytokine production in vitro and exhibits synergism when combined with IL-12 in terms of enhanced IFN- γ production (Micallef et al., 1996).

After the murine form was cloned, the human cDNA sequence for IL-18 was reported in 1996 (Okamura et al., 1995).

5 By cloning IL-18 from affected tissues and studying IL-18 gene expression, a close association of this cytokine with an autoimmune disease was found. The non-obese diabetic (NOD) mouse spontaneously develops autoimmune insulinitis and diabetes, which can be accelerated and synchronized by a single injection of cyclophosphamide. IL-18 mRNA was demonstrated by reverse transcriptase PCR in
10 NOD mouse pancreas during early stages of insulinitis. Levels of IL-18 mRNA increased rapidly after cyclophosphamide treatment and preceded a rise in IFN- γ mRNA, and subsequently diabetes. Interestingly, these kinetics mimic that of IL-12-p40 mRNA, resulting in a close correlation of individual mRNA levels. Cloning of the IL-18 cDNA from pancreas RNA followed by sequencing revealed identity with the IL-18 sequence
15 cloned from Kupffer cells and in vivo pre-activated macrophages. Also NOD mouse macrophages responded to cyclophosphamide with IL-18 gene expression while macrophages from Balb/c mice treated in parallel did not. Therefore, IL-18 expression is abnormally regulated in autoimmune NOD mice and closely associated with diabetes development (Rothe et al., 1997).

20 IL-18 plays a potential role in immunoregulation or in inflammation by augmenting the functional activity of Fas ligand on Th1 cells (Conti et al., 1997). IL-18 is also expressed in the adrenal cortex and therefore might be a secreted neuro-immunomodulator, playing an important role in orchestrating the immune system following a stressful experience (Chater, 1986).

25 In vivo, IL-18 is formed by cleavage of pro-IL-18, and its endogenous activity appears to account for IFN- γ production in *P. acnes* and LPS-mediated lethality. Mature IL-18 is produced from its precursor by the IL-1 β converting enzyme (IL-1 β -converting enzyme, ICE, caspase-1).

The IL-18 receptor consists of at least two components, co-operating in ligand
30 binding. High- and low-affinity binding sites for IL-18 were found in murine IL-12 stimulated T cells (Yoshimoto et al., 1998), suggesting a multiple chain receptor complex. Two receptor subunits have been identified so far, both belonging to the IL-1

receptor family (Parnet et al., 1996). The signal transduction of IL-18 involves activation of NF- κ B (DiDonato et al., 1997).

5 Recently, a soluble protein having a high affinity for IL-18 has been isolated from human urine, and the human and mouse cDNAs were described (Novick et al., 1999; WO 99/09063). The protein has been designated IL-18 binding protein (IL-18BP).

10 IL-18BP is not the extracellular domain of one of the known IL18 receptors, but a secreted, naturally circulating protein. It belongs to a novel family of secreted proteins. The family further includes several Poxvirus-encoded proteins which have a high homology to IL-18BP (Novick et al., 1999). IL-18BP is constitutively expressed in the spleen, belongs to the immunoglobulin superfamily, and has limited homology to the IL-1 type II receptor. Its gene was localized on human chromosome 11q13, and no exon coding for a transmembrane domain was found in an 8.3 kb genomic sequence
15 (Novick et al., 1999).

Four human and two mouse isoforms of IL-18BP, resulting from mRNA splicing and found in various cDNA libraries and have been expressed, purified, and assessed for binding and neutralization of IL-18 biological activities (Kim et al., 2000). Human IL-18BP *isoform a* (IL-18BP_a) exhibited the greatest affinity for IL-18 with a rapid on-rate, a slow off-rate, and a dissociation constant (K(d)) of 399 pM. IL-18BP_c shares the Ig domain of IL-18BP_a except for the 29 C-terminal amino acids; the K(d) of IL-18BP_c is 10-fold less (2.94 nM). Nevertheless, IL-18BP_a and IL-18BP_c neutralize IL-18 >95% at a molar excess of two. IL-18BP_b and IL-18BP_d isoforms lack a complete Ig domain and lack the ability to bind or neutralize IL-18. Murine IL-18BP_c and IL-18BP_d isoforms, 25 possessing the identical Ig domain, also neutralize >95% murine IL-18 at a molar excess of two. However, murine IL-18BP_d, which shares a common C-terminal motif with human IL-18BP_a, also neutralizes human IL-18. Molecular modeling identified a large mixed electrostatic and hydrophobic binding site in the Ig domain of IL-18BP, which could account for its high affinity binding to the ligand (Kim et al., 2000).

30 A beneficial effect of IL-18BP in several diseases has been described. Examples of such diseases treatable with IL-18BP are: tumor metastasis (WO 01/07480), arthritis, inflammatory bowel disease and liver injury (WO 01/62285), heart

disease (WO 02/060479), traumatic brain injury (WO 02/096456), sepsis (WO 02/092008), atherosclerosis (WO 01/85201), hypersensitivity disorder (WO03/033015).

An IL-18 binding domain of a viral homologue of human IL-18BP has been described in the literature (Xiang and Moss, 2003), showing that a furin cleaved form of
5 the IL-18 binding protein of Mollusum contagiosum virus has IL-18 binding properties. In addition to this, several point mutations were introduced into the viral homologue of IL-18BP and tested in vitro for IL-18 binding in order to define the biologically active portions of the protein (Xiang and Moss, 2001).

However, biologically active fragments of the human IL-18BP *isoform a* have not
10 been identified and characterized so far.

SUMMARY OF THE INVENTION

The present invention is based on the finding that during production of recombinant human IL-18BP *isoform a*, truncated forms can be found that retain the
15 biological activity of IL-18BP, as measured in an in vitro bioassay. The invention therefore relates to variants of the IL-18BP having amino acid sequences of SEQ ID NO: 2, 3, 4, 5, 6 or 7, and to variants lacking the C-terminal amino acid residue of these sequences. Such variants represent active variants of the mature IL-18BP.

The invention further relates to nucleic acid molecules coding for such IL-18BP
20 variants, to pharmaceutical compositions comprising these variants, and to their use for the treatment and/or prevention of IL-18 mediated disorders.

In a further aspect, the invention relates to the use of an expression vector comprising the coding sequence of an IL-18BP variant for the treatment and/or prevention of IL-18 mediated diseases.

25 The invention further relates to processes of production of the IL-18BP variants of the invention.

BRIEF DESCRIPTION OF THE DRAWINGS

Fig. 1 shows the sequence of full-length IL-18BP *isoform a*. Putative N-glycosylation
30 sites are labeled as N*. Arrows mark the N-termini of the six IL-18BP variants of the invention.

Fig. 2 shows a silver stained SDS-PAGE gel (A) and the corresponding western blot (B) of an IL-18BP preparation containing IL-18BP variants of the invention. The lanes were loaded as follows:

5

Fig. 2 A:

1. MW marker
2. r-hIL-18BP CT20 (2 µg)
3. r-hIL-18BP CT20 (2 µg)
4. r-hIL-18BP CT20 (2 µg)
5. ST1P01/r-hIL-18BP

Fig. 2B

1. MW marker
2. r-hIL-18BP CT20 (200 ng)
3. ST1P01/r-hIL-18BP (200 ng)
4. MW marker

10

Fig. 3. shows the SE-HPLC profile as well as silver stained SDS-PAGE gel (A) and the corresponding western blot (B) of the two peaks obtained in HPLC as compared to a standard preparation of pure full-length IL-18BP *isoform a*.

15

DESCRIPTION OF THE INVENTION

The present invention is based on the finding that variants of IL-18BP could be identified during recombinant production of human recombinant IL-18BP *isoform a*. These variants were characterized and it was found that they represent defined N- and C-terminally truncated fragments of full-length IL-18BP *isoform a*. Definition of the N-glycosylation pattern of recombinant IL-18BP could be achieved in the frame of the present invention, leading to a new variant of full-length IL-18BP.

20

Surprisingly, all variants of IL-18BP displayed a biological activity comparable to full-length IL-18BP in an *in vitro* bioassay.

Therefore, in a first aspect, the invention relates to a new IL-18 binding protein (IL-18BP) comprising an amino acid sequence selected from SEQ ID NO: 2, 3, 4, 5, 6 or 7, but not SEQ ID NO: 1, or functional derivatives, fusion proteins or salts thereof. The invention thus relates to active fragments of the IL-18BP containing defined

25

portions of full-length IL-18BP, but not the full-length sequence of IL-18BP *isoform a*, which is depicted in SEQ ID NO: 1.

In a preferred embodiment, the IL-18BP consists of an amino acid sequence selected from SEQ ID NO: 2, 3, 4, 5, 6 or 7.

5 In the frame of the present invention, it has further been found that there is a C-terminal heterogeneity of the IL-18BP variants in that species lacking the very C-terminal residue can be detected to some extent.

Therefore, in a preferred embodiment, the invention relates to an IL-18 binding protein (IL-18BP) comprising or consisting of an amino acid sequence selected from
10 SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, or SEQ ID NO: 7, but not comprising or consisting of SEQ ID NO: 1, less the C-terminal glycine residue, or a functional derivative, fusion protein or salt thereof.

In the frame of the present invention, variants of IL-18BP have been identified, in which an internal clipping of IL-18BP has occurred.

15 In a second aspect, the invention relates to an IL-18BP comprising a first polypeptide consisting of amino acids 1 to 30 of SEQ ID NO: 1 and a second polypeptide consisting of amino acids 31 to 164 or of amino acids 31 to 163, wherein the first and second polypeptide are linked by a disulfide bond.

In a third aspect, the IL-18BP comprises a first polypeptide consisting of amino
20 acids 15 to 30 of SEQ ID NO: 1 and a second polypeptide consisting of amino acids 31 to 164 or of amino acids 31 to 163, wherein the first and second polypeptide are linked by a disulfide bond.

The IL-18BPs may be unglycosylated or glycosylated. Preferably, the IL-18BPs of the invention are N-glycosylated at asparagine residues Asn 49, Asn 73 and Asn 117
25 (numbering according to Fig. 1).

During the experiments leading to the present invention, it has been found for the first time that recombinantly produced full-length IL-18 binding protein *isoform a* is not glycosylated at all putative N-glycosylation sites, but only at three defined Asparagine residues, which are Asn 49, 73 and 117. Therefore, the invention further
30 relates to an IL-18BP having the amino acid sequence of SEQ ID NO: 1, wherein the protein is N-glycosylated at Asn 49, Asn 73 and Asn 117, as well as to functional derivatives, fusion proteins or salts thereof.

In a preferred embodiment, the IL-18BP has the amino acid sequence of SEQ ID NO: 1 less the C-terminal glycine residue.

The sequence of full length human IL-18BP and its splice variants/isoform are disclosed e.g. from WO99/09063, or from Novick et al., 1999, as well as in Kim et al.,
5 2000. SEQ ID NO: 1 represents the amino acid sequence of mature full-length IL-18BP *isoform a*.

In the following, the proteins of the invention may be generally designated "IL-18BP", "IL-18BPs" or "IL-18BP(s) of the invention". These terms, as used therein, encompass all IL-18BP variants described in the frame of the present invention.

10 The proteins according to the present invention may be derived from natural sources, such as urine, or they may preferably be produced recombinantly. Recombinant expression may be carried out in prokaryotic expression systems like E. coli, or in eukaryotic, and preferably in mammalian, expression systems. They may also preferably be produced in human expression systems. Established cell lines such as
15 the Chinese hamster ovary cell line (CHO) or the human embryonic kidney cell line 293 may be especially useful for production of the IL-18BP variants of the present invention.

Further variants within the scope of the present invention may be proteins having conservative amino acid substitutions of the sequences depicted in Fig. 1 or the
20 annexed sequence listing. These variants may be prepared by known synthesis and/or by site-directed mutagenesis techniques, or any other known technique suitable therefor.

Conservative amino acid substitutions of IL-18BP polypeptides, may include synonymous amino acids within a group which have sufficiently similar physicochemical
25 properties that substitution between members of the group will preserve the biological function of the molecule (Grantham, 1974). It is clear that insertions and deletions of amino acids may also be made in the above-defined sequences without altering their function, particularly if the insertions or deletions only involve a few amino acids, e.g., under thirty, and preferably under ten, and do not remove or displace amino acids
30 which are critical to a functional conformation, e.g., cysteine residues. Proteins and muteins produced by such deletions and/or insertions come within the purview of the present invention.

Preferably, the synonymous amino acid groups are those defined in Table 1.
More preferably, the synonymous amino acid groups are those defined in Table 2; and
most preferably the synonymous amino acid groups are those defined in Table 3.

5

TABLE 1

Preferred Groups of Synonymous Amino Acids

	Amino Acid	Synonymous Group
	Ser	Ser, Thr, Gly, Asn
	Arg	Arg, Gln, Lys, Glu, His
10	Leu	Ile, Phe, Tyr, Met, Val, Leu
	Pro	Gly, Ala, Thr, Pro
	Thr	Pro, Ser, Ala, Gly, His, Gln, Thr
	Ala	Gly, Thr, Pro, Ala
	Val	Met, Tyr, Phe, Ile, Leu, Val
15	Gly	Ala, Thr, Pro, Ser, Gly
	Ile	Met, Tyr, Phe, Val, Leu, Ile
	Phe	Trp, Met, Tyr, Ile, Val, Leu, Phe
	Tyr	Trp, Met, Phe, Ile, Val, Leu, Tyr
	Cys	Ser, Thr, Cys
20	His	Glu, Lys, Gln, Thr, Arg, His
	Gln	Glu, Lys, Asn, His, Thr, Arg, Gln
	Asn	Gln, Asp, Ser, Asn
	Lys	Glu, Gln, His, Arg, Lys
	Asp	Glu, Asn, Asp
25	Glu	Asp, Lys, Asn, Gln, His, Arg, Glu
	Met	Phe, Ile, Val, Leu, Met
	Trp	Trp

TABLE 2

30

More Preferred Groups of Synonymous Amino Acids

Amino Acid	Synonymous Group
Ser	Ser
Arg	His, Lys, Arg

	Leu	Leu, Ile, Phe, Met
	Pro	Ala, Pro
	Thr	Thr
	Ala	Pro, Ala
5	Val	Val, Met, Ile
	Gly	Gly
	Ile	Ile, Met, Phe, Val, Leu
	Phe	Met, Tyr, Ile, Leu, Phe
	Tyr	Phe, Tyr
10	Cys	Cys, Ser
	His	His, Gln, Arg
	Gln	Glu, Gln, His
	Asn	Asp, Asn
	Lys	Lys, Arg
15	Asp	Asp, Asn
	Glu	Glu, Gln
	Met	Met, Phe, Ile, Val, Leu
	Trp	Trp

20

TABLE 3

Most Preferred Groups of Synonymous Amino Acids

	Amino Acid	Synonymous Group
	Ser	Ser
	Arg	Arg
25	Leu	Leu, Ile, Met
	Pro	Pro
	Thr	Thr
	Ala	Ala
	Val	Val
30	Gly	Gly
	Ile	Ile, Met, Leu
	Phe	Phe
	Tyr	Tyr

	Cys	Cys, Ser
	His	His
	Gln	Gln
	Asn	Asn
5	Lys	Lys
	Asp	Asp
	Glu	Glu
	Met	Met, Ile, Leu
	Trp	Met

10

It is understood that minor changes in the amino acid sequence of the IL-18BP variants are within the scope of the invention, having a sequence of amino acids sufficiently duplicative of that of an IL-18BP variant described herein, such as to have a comparable activity to IL-18BP. One activity of IL-18BP is its capability of binding IL-18.

15 Thus, it can be determined whether any given variant has substantially the same activity as IL-18BP by means of routine experimentation comprising subjecting such a mutein, e.g., to a simple sandwich competition assay to determine whether or not it binds to an appropriately labeled IL-18, such as radio-immunoassay or ELISA assay. A further meaningful assay describing IL-18BP activity is the bioassay described in the

20 example below.

Examples of production of amino acid substitutions in proteins which can be used for obtaining variants of IL-18BP polypeptides or proteins for use in the present invention include any known method steps, such as presented in US patents 4,959,314, 4,588,585 and 4,737,462, to Mark et al; 5,116,943 to Koths et al., 4,965,195

25 to Namen et al; 4,879,111 to Chong et al; and 5,017,691 to Lee et al; and lysine substituted proteins presented in US patent No. 4,904,584 (Shaw et al).

In an embodiment of the invention, the IL-18BP variants are fused proteins.

The term "fused protein" refers to a polypeptide comprising an IL-18BP of the invention, fused with another protein, which, e.g., has an extended residence time in

30 body fluids. An IL-18BP may thus be fused to another protein, polypeptide or the like, e.g., an immunoglobulin or a fragment thereof.

In a preferred embodiment of the invention, the IL -18BP of the invention comprises an immunoglobulin fusion, i.e. it is a fused protein comprising all or part of

an IL-18BP of the invention, which is fused to all or a portion of an immunoglobulin. Methods for making immunoglobulin fusion proteins are well known in the art, such as the ones described in WO 01/03737, for example. The person skilled in the art will understand that the resulting fusion protein of the invention retains the biological activity of IL-18BP, in particular the binding to IL-18. The fusion may be direct, or via a short linker peptide which can be as short as 1 to 3 amino acid residues in length or longer, for example, 13 amino acid residues in length. Said linker may be a tripeptide of the sequence E-F-M (Glu-Phe-Met), for example, or a 13-amino acid linker sequence comprising Glu-Phe-Gly-Ala-Gly-Leu-Val-Leu-Gly-Gly-Gln-Phe-Met introduced between the IL-18BP sequence and the immunoglobulin sequence. The resulting fusion protein has improved properties, such as an extended residence time in body fluids (half-life), increased specific activity, increased expression level, or the purification of the fusion protein may be facilitated.

In a preferred embodiment, IL-18BP is fused to the constant region of an Ig molecule. Preferably, it is fused to heavy chain regions, like the CH2 and CH3 domains of human IgG1 or IgG3, for example. The generation of specific fusion proteins comprising IL-18BP and a portion of an immunoglobulin are described in example 11 of WO 99/09063, for example. Other isoforms of Ig molecules are also suitable for the generation of fusion proteins according to the present invention, such as isoforms IgG₂ or IgG₄, or other Ig classes, like IgM or IgA, for example. Fusion proteins may be monomeric or multimeric, hetero- or homomultimeric.

The invention further relates to a process for production of an IL-18BP fused protein comprising preparing a DNA construct that encodes an IL-18BP of the invention ligated to a nucleic acid encoding a second polypeptide, wherein upon expression, said DNA construct encodes a fusion protein comprising the IL-18BP of the invention fused to the second polypeptide.

Preferably, the second polypeptide is a portion of an immunoglobulin, more preferably the Fc portion of an immunoglobulin.

In a further embodiment, the IL-18BP variants are functional derivatives.

"Functional derivatives" as used herein cover derivatives of IL-18BP variants or their fused proteins, which may be prepared from the functional groups which occur as side chains on the residues or the N- or C-terminal groups, by means known in the art, and are included in the invention as long as they remain pharmaceutically acceptable,

i.e. they do not destroy the activity of the protein which is substantially similar to the activity of IL-18BP, or viral IL-18BPs, and do not confer toxic properties on compositions containing it. Functional derivatives of IL-18BP may be conjugated to polymers in order to improve the properties of the protein, such as the stability, half-life, bioavailability, tolerance by the human body, or immunogenicity. To achieve this goal, IL18-BP may be linked e.g. to Polyethyenglycol (PEG). PEGylation may be carried out by known methods, described in WO 92/13095, for example.

Derivatives may also, for example, include aliphatic esters of the carboxyl groups, amides of the carboxyl groups by reaction with ammonia or with primary or secondary amines, N-acyl derivatives of free amino groups of the amino acid residues formed with acyl moieties (e.g. alkanoyl or carbocyclic aroyl groups) or O-acyl derivatives of free hydroxyl groups (for example that of seryl or threonyl residues) formed with acyl moieties.

The invention further relates to a process for production of an IL-18BP derivative of the invention comprising chemically modifying an IL-18BP of the invention to include at least one derivative moiety. Preferably, the moiety is a polyethylene glycol moiety.

Yet a further embodiment of the invention relates to salts of the IL-18BP variants.

The term "salts" herein refers to both salts of carboxyl groups and to acid addition salts of amino groups of IL-18BP variant molecule, or analogs thereof. Salts of a carboxyl group may be formed by means known in the art and include inorganic salts, for example, sodium, calcium, ammonium, ferric or zinc salts, and the like, and salts with organic bases as those formed, for example, with amines, such as triethanolamine, arginine or lysine, piperidine, procaine and the like. Acid addition salts include, for example, salts with mineral acids, such as, for example, hydrochloric acid or sulfuric acid, and salts with organic acids, such as, for example, acetic acid or oxalic acid. Of course, any such salts must retain the biological activity of the IL-18BP relevant to the present invention, such as inhibition of IFN-gamma induction in the bioassay described in the examples below.

In a further aspect, the invention relates to a nucleic acid coding for an IL-18BP of the invention. Such coding sequence may easily be deduced from the amino acid sequences depicted in Fig. 1 or the annexed sequence listing. The person skilled in the

art will appreciate that many more nucleic acid sequences coding for the IL-18BPs of the invention can be conceived due to the degeneracy of the genetic code.

In yet a further aspect, the invention relates to a host cell comprising the nucleic acid of the invention. Such a host cell may be either prokaryotic or eukaryotic, preferably mammalian, more preferably a host cell suitable for recombinant expression
5 of therapeutic proteins such as Chinese hamster ovary cells (CHO) or human cells.

The invention further relates to a process for production of an IL-18BP of the invention comprising the step of culturing a host cell according to the invention under conditions suitable for expression of said IL-18BP.

10 The process for production of an IL-18BP may also comprise the step of isolating the IL-18BP from the cell culture supernatant of a host cell of the invention.

In another aspect, the invention relates to a composition comprising an IL-18BP in accordance with the present invention. Preferably, it is a pharmaceutical composition. Optionally, the pharmaceutical composition further comprises
15 pharmaceutically acceptable surfactants, excipients, carriers, diluents and vehicles.

The definition of "pharmaceutically acceptable" is meant to encompass any carrier, which does not interfere with effectiveness of the biological activity of the active ingredient and that is not toxic to the host to which it is administered. For example, for parenteral administration, the active protein(s) may be formulated in a unit dosage form
20 for injection in vehicles such as saline, dextrose solution, serum albumin and Ringer's solution.

The active ingredients of the pharmaceutical composition according to the invention can be administered to an individual in a variety of ways. The routes of administration include intradermal, transdermal (e.g. in slow release formulations),
25 intramuscular, intraperitoneal, intravenous, subcutaneous, oral, intracranial, epidural, topical, rectal, and intranasal routes.

Preferred administration routes of the invention are the subcutaneous and the intramuscular route.

Any other therapeutically efficacious route of administration can be used, for
30 example absorption through epithelial or endothelial tissues, or by gene therapy wherein a DNA molecule encoding the active agent is administered to the patient (e.g. via a vector), which causes the active agent to be expressed and secreted in vivo. If an

expression vector comprising the coding sequence of IL-18BP(s) of the invention is to be administered, it may e.g. be injected intramuscularly as naked DNA.

For parenteral (e.g. intravenous, subcutaneous, intramuscular) administration, the active protein(s) can be formulated as a solution, suspension, emulsion or lyophilized powder in association with a pharmaceutically acceptable parenteral vehicle (e.g. water, saline, dextrose solution) and additives that maintain isotonicity (e.g. mannitol) or chemical stability (e.g. preservatives and buffers). The formulation is sterilized by commonly used techniques.

The bioavailability of the active protein(s) according to the invention can also be ameliorated by using conjugation procedures which increase the half-life of the molecule in the human body, for example linking the molecule to polyethylene glycol, as described in the PCT Patent Application WO 92/13095.

The therapeutically effective amounts of the active protein(s) will be a function of many variables, including the type of IL-18BP use, their affinity for IL-18, any residual cytotoxic activity exhibited by the IL-18BP(s), the route of administration, the clinical condition of the patient (including the desirability of maintaining a non-toxic level of endogenous IL-18 activity).

A "therapeutically effective amount" is such that when administered, the IL-18BP variant results in inhibition of the biological activity of IL-18. The dosage administered, as single or multiple doses, to an individual will vary depending upon a variety of factors, including IL-18BP variant pharmacokinetic properties, the route of administration, patient conditions and characteristics (sex, age, body weight, health, size), extent of symptoms, concurrent treatments, frequency of treatment and the effect desired. Adjustment and manipulation of established dosage ranges are well within the ability of those skilled in the art, as well as in vitro and in vivo methods of determining the inhibition of IL-18 in an individual.

In a preferred embodiment of the present invention, the IL-18BP variant is used in an amount of about 0.001 to 1000 mg/kg of body weight, or about 0.001 to 100 mg/kg of body weight or about 0.01 to 10 mg/kg of body weight or about 0.1 to 5 mg/kg or about 1 to 3 mg/kg of body weight.

The frequency of administration may be daily or every other day. It may also be three times per week or once per week.

The doses administered may always be the same or vary, depending on the patient's needs. The doses are usually given in divided doses or in sustained release form effective to obtain the desired results. Second or subsequent administrations can be performed at a dosage which is the same, less than or greater than the initial or
5 previous dose administered to the individual. A second or subsequent administration can be administered during or prior to onset of the disease.

According to the invention, the IL-18BP variant can be administered prophylactically or therapeutically to an individual prior to, simultaneously or sequentially with other therapeutic regimens or agents (e.g. multiple drug regimens), in
10 a therapeutically effective amount, in particular with an interferon and/or a TNF inhibitor. Active agents that are administered simultaneously with other therapeutic agents can be administered in the same or different compositions.

In a further aspect, the invention relates to the use of an IL-18BP of the invention for the preparation of a medicament for treatment and/or prevention of an IL-
15 18 mediated disease or disorder. IL-18 mediated diseases are known in the art (reviewed e.g. by Gracie et al., 2003).

In a preferred embodiment, the disease to be treated or prevented by the IL-18BP variant of the invention is selected from psoriasis, arthritis, in particular rheumatoid arthritis, inflammatory bowel disease, in particular Crohn's disease, liver injury,
20 atherosclerosis, sepsis, myocardial infarction, traumatic brain injury, allergy, peripheral vascular disease, multiple sclerosis, tumor metastasis.

For detailed description and definition of these diseases, it is particularly referred to the following published patent applications which are fully incorporated by reference herein: WO 99/09063, WO 01/07480, WO 01/62285, WO 02/060479, WO
25 02/096456, WO 02/092008, WO 03/013577.

Interferons are predominantly known for inhibitory effects on viral replication and cellular proliferation. Interferon- γ , for example, plays an important role in promoting immune and inflammatory responses. Interferon β (IFN- β , an interferon type I), is said to play an anti-inflammatory role.

30 The invention therefore also relates to the use of a combination of an IL-18BP of the invention and an interferon in the manufacture of a medicament for the treatment of an IL-18 mediated disease.

Interferons may also be conjugated to polymers in order to improve the stability of the proteins. A conjugate between Interferon β and the polyol Polyethyenglycol (PEG) has been described in WO99/55377, for instance.

In another preferred embodiment of the invention, the interferon is Interferon- β (IFN- β), and more preferably IFN- β 1a.

The IL-18BP of the invention is preferably used simultaneously, sequentially, or separately with the interferon.

In yet a further embodiment of the invention, an IL-18BP of the invention is used in combination with a TNF antagonist. TNF antagonists exert their activity in several ways. First, antagonists can bind to or sequester the TNF molecule itself with sufficient affinity and specificity to partially or substantially neutralize the TNF epitope or epitopes responsible for TNF receptor binding (hereinafter termed "sequestering antagonists"). A sequestering antagonist may be, for example, an antibody directed against TNF.

Alternatively, TNF antagonists can inhibit the TNF signaling pathway activated by the cell surface receptor after TNF binding (hereinafter termed "signaling antagonists"). Both groups of antagonists are useful, either alone or together, in combination with an IL-18BP variant, in the therapy of hypersensitivity disorders.

TNF antagonists are easily identified and evaluated by routine screening of candidates for their effect on the activity of native TNF on susceptible cell lines in vitro, for example human B cells, in which TNF causes proliferation and immunoglobulin secretion. The assay contains TNF formulation at varying dilutions of candidate antagonist, e.g. from 0.1 to 100 times the molar amount of TNF used in the assay, and controls with no TNF or only antagonist (Tucci et al., 1992).

Sequestering antagonists are the preferred TNF antagonists to be used according to the present invention. Amongst sequestering antagonists, those polypeptides that bind TNF with high affinity and possess low immunogenicity are preferred. Soluble TNF receptor molecules and neutralizing antibodies to TNF are particularly preferred. For example, soluble TNF-RI (also called p55) and TNF-RII (also called p75) are useful in the present invention. Truncated forms of these receptors, comprising the extracellular domains of the receptors or functional portions thereof, are more particularly preferred antagonists according to the present invention. Soluble TNF

type-I and type-II receptors are described in European Patents EP 308 378, EP 398 327 and EP 433 900, for example.

These truncated, soluble TNF receptors are soluble and have been detected in urine and serum as TNF inhibitory binding proteins, called TBPI and TBPII, respectively
5 (Engelmann et al., 1990). The simultaneous, sequential, or separate use of the IL-18BP variant with the TNF antagonist and /or an Interferon is preferred, according to the invention.

According to the invention, TBP I and TBPII are preferred TNF antagonists to be used in combination with an IL-18BP variant of the invention. Derivatives, fragments,
10 regions and biologically active portions of the receptor molecules functionally resemble the receptor molecules that can also be used in the present invention. Such biologically active equivalent or derivative of the receptor molecule refers to the portion of the polypeptide, or of the sequence encoding the receptor molecule, that is of sufficient size and able to bind TNF with such an affinity that the interaction with the membrane-
15 bound TNF receptor is inhibited or blocked.

The invention further relates to the use of an expression vector comprising the coding sequence of an IL-18BP of the invention in the preparation of a medicament for the prevention and/or treatment of IL-18 mediated disorders. Thus, a gene therapy approach is considered in order to deliver the IL-18BP variant to the site where it
20 required. In order to treat and/or prevent a hypersensitivity disorder, the gene therapy vector comprising the sequence of an IL-18BP variant production and/or action may be injected directly into the diseased tissue, for example, thus avoiding problems involved in systemic administration of gene therapy vectors, like dilution of the vectors, reaching and targeting of the target cells or tissues, and of side effects.

25 The invention further relates to the use of a cell that has been genetically modified to produce an IL-18BP of the invention in the manufacture of a medicament for the treatment and/or prevention of an IL-18 mediated disease.

The invention further relates to a method for the preparation of a pharmaceutical composition comprising admixing an effective amount of an IL-18BP
30 variant and/or an interferon and/or a TNF antagonist with a pharmaceutically acceptable carrier.

The invention further relates to a method of treatment of IL-18 mediated disease, comprising administering a pharmaceutically effective amount of an IL-18BP variant to a patient in need thereof.

5 Having now fully described this invention, it will be appreciated by those skilled in the art that the same can be performed within a wide range of equivalent parameters, concentrations and conditions without departing from the spirit and scope of the invention and without undue experimentation.

10 While this invention has been described in connection with specific embodiments thereof, it will be understood that it is capable of further modifications. This application is intended to cover any variations, uses or adaptations of the invention following, in general, the principles of the invention and including such departures from the present disclosure as come within known or customary practice within the art to which the invention pertains and as may be applied to the essential features hereinbefore set forth as follows in the scope of
15 the appended claims.

 All references cited herein, including journal articles or abstracts, published or unpublished U.S. or foreign patent application, issued U.S. or foreign patents or any other references, are entirely incorporated by reference herein, including all data, tables, figures and text presented in the cited references. Additionally, the entire contents of the references
20 cited within the references cited herein are also entirely incorporated by reference.

 Reference to known method steps, conventional methods steps, known methods or conventional methods is not any way an admission that any aspect, description or embodiment of the present invention is disclosed, taught or suggested in the relevant art.

 The foregoing description of the specific embodiments will so fully reveal the general
25 nature of the invention that others can, by applying knowledge within the skill of the art (including the contents of the references cited herein), readily modify and/or adapt for various application such specific embodiments, without undue experimentation, without departing from the general concept of the present invention. Therefore, such adaptations and modifications are intended to be within the meaning an range of equivalents of the
30 disclosed embodiments, based on the teaching and guidance presented herein. It is to be understood that the phraseology or terminology herein is for the purpose of description and not of limitation, such that the terminology or phraseology of the present specification is to

be interpreted by the skilled artisan in light of the teachings and guidance presented herein,
in combination with the knowledge of one of ordinary skill in the art.

5

EXAMPLE

IDENTIFICATION OF IL-18BP VARIANTS

10

MATERIALS and METHODS

Materials and equipment

	96 well microtiter plate photometer MCC 349 or EX	Labsystem
	Analytical balance mod. AG145	Mettler-Toledo
15	Aquapore RP300 30 x 4.6 mm cartridge cod. 0711-0055	Brownlee
	Automated sequencer mod. Procise 494	Applied Biosystem
	Automatic pipettes (P1000, P200, P100, P20)	Gilson
	Cell Coulter	Counter -Z1
	CO ₂ incubator	Heraeus
20	Excel software	
	Freezer -20°C ± 5°C	Angelantoni
	Freezer -80°C ± 10°C	Angelantoni
	Graph Pad Prism Software	
	HPLC mod. Alliance 2690	Waters
25	HPLC-pump mod. 600S with column heater	Waters
	Integrator D2500	Merck
	Laminar Flow Hood	Flow Laboratories
	MALDI-ToF mod. Voyager DE-Pro	Perseptive Biosystem
	Mass Spectrometer mod. ZQ	Waters Micromass
30	Multiphor II	Pharmacia
	Multitemp II	Pharmacia or equivalent
	Personal computer	CompaQ
	pH meter MA235	Mettler or equivalent

	pH-meter mod MP225	Mettler-Toledo
	Power supply EPS 3501 XL	Pharmacia or equivalent
	Refrigerator +5°C ± 3°C	Angelantoni
	Scanner AGFA Arcus II	Agfa or equivalent
5	Separation module 2690 Alliance	Waters
	Software Agfa Fotolook v.3.0	Agfa or equivalent
	Software Millennium ³² version 3.20	Waters
	Software Phoretix 1D	Phoretix or equivalent
	Software Picture Publisher v.8	Micrografx or equivalent
10	Spectrolinker XL 1000 Cross Linker (UV source)	Spectronics Corporation
	Statgraphics Plus Symmetry C18 3.5 µm 75 x 4,6 mm column cod. WAT066224	Waters
	Technical balance mod PEG2002	Mettler-Toledo
	UV detector 2487	Waters
	UV detector mod. 2487	Waters
15	UV detector mod. 996	Waters

Chemicals

	Dithiothreitol (DTT) cod. D5545	Sigma
	Tris cod.1.08382	Merck
20	EDTA cod.1.08418	Merck
	Acetonitrile (ACN) cod. 1.00030	Merck
	Ammonium bicarbonate cod. 1.01131	Merck
	Ammonia 25% cod. 1.05432	Merck
	Calcium chloride 2 H ₂ O cod. I3381	Sigma
25	Endoproteinase Bovine Trypsin, Modified, sequencing grade cod. 1418-025	Roche
	Neuraminidase (Sialidase) cod. 1080-725	Roche
	Water (H ₂ O) MilliQ Grade	Millipore
	Trifluoroacetic acid (TFA) cod. 9470	Baker
	Acetic acid glacial cod. 00063	Merck
30	Sodium Hydroxide cod. 7067	Baker
	Iodoacetic acid cod. I2512	Sigma
	Sodium acetate 3M pH 5.5 cod. 400471	Applied Biosystem
	Hydrochloric acid 37% cod.1.000314	Merck

	β -Mercaptoethanol cod. M 6250	Sigma
	Diethylether cod. cod 447521	Carlo Erba
	Guanidine cod. N24115	Pierce
	NaCsl cod. 700000889-2	ULTRA Scientific
5	Nitrogen UPP	Caracciolo
	Methanol gradient grade cod 1.06007	Merck
	Eppendorf 1.5 ml	Eppendorf
	Water (H ₂ O) purified by Modulab 2020™	Continental
	Acetonitrile (HPLC grade) code 1.00030	Merck
10	o-Phosphoric acid (H ₃ PO ₄) 85% code 1.00573	Merck
	Sodium sulfate (Na ₂ SO ₄) code 1.06649	Merck
	Column TSK G2000 SW _{XL} 7.8x300 code 08540	TosoHaas
	Goat anti mouse IgG HRP conjugated cod. 170-6516	BioRad
	Monoclonal antibody anti r-hIL-18BP clone 582.10	IPL
15	ExcelGel SDS buffer strips cod. 17-1342-01	Pharmacia
	ExcelGel SDS Homogeneous 12.5% cod. 80-1261-01	Pharmacia
	Hyperfilm ECL 18 x 24 cm cod. RPN 2103	Pharmacia-Biotech
	Kit ECL cod. RPN2106	Pharmacia-Biotech
	Kit Silver PlusOne cod. 17-1150-01	Pharmacia
20	Nitrocellulose membrane 0.2mm cod. BA-83	Schleicher & Schuell
	I-Block cod. AI 300	Tropix
	Interim Reference Material ST1P01/r-hIL-18BP	IFS
	Molecular weight marker (97-14 kDa) cod. 17-0446-01	Pharmacia
	Tween 20	Merck
25	Phosphate buffered saline (PBS). with calcium and magnesium ions.	Sigma
	96 wells plate	Falcon
	96 wells microtiter plate	Maxi Sorp Nunc
	IMDM	GIBCO
	2-Mercaptoethanol	Sigma
30	Penicilin/Streptomycin	Gibco
	Foetal Bovine Serum (FBS)	GIBCO
	Human IFN- γ Immunoassay Kit – DUO Set ELISA Development System	R&D Systems
	Bovine Serum Albumin (BSA)	Sigma

Substrate solution
Sulphuric Acid (H₂SO₄)

R&D Systems
Merck

Biologicals

- 5 Human acute myelogenous leukemia cell line KG-1 In house
Recombinant human Tumor Necrosis Factor- α (TNF- α) R&D Systems
Recombinant human Interleukin-18 (r-hIL-18) Produced in house
Recombinant human Interleukin-18 Binding Protein-(r-hIL-18BP) produced in house
(46.39 mg/ml by Amino Acid Analysis)

10

METHODS

PEPTIDE MAPPING BY TRYPSIN

The mapping was carried out according to standard protocols, outlined below.

15

TREATMENT WITH NEURAMINIDASE

About 150 μ g of dried r-hIL-18BP was dissolved with 200 μ L of 0.2M Ammonium Acetate 16mM Calcium Chloride pH 5.5 buffer and 100mIU of Sialidase. The reaction was performed at 37 \pm 1 $^{\circ}$ C for 1 hour. Then the protein was dried in Speed-Vacuum.

- 20 After desiccation the protein was reduced and alkylated as described below.

REDUCTION AND ALKYLATION

- Dissolved with 200 μ L of 0.5M Tris-Cl 2mM EDTA pH 8,5 \pm 0.05 6M Guanidine 11mg/mL dithiotreitol under nitrogen atmosphere. The reaction was performed at room
25 temperature for 1 hour. Then has been added 25 μ L of 250mg/mL iodoacetic acid under nitrogen atmosphere. The mixture was incubated in the dark at 37 \pm 1 $^{\circ}$ C for 45 minutes and then stopped by adding 200 μ L of 0.1% aqueous TFA and 20 μ L β -mercaptoethanol under nitrogen atmosphere. The reaction was incubated at room temperature for 15 minutes.

30

PURIFICATION PROCEDURE

After reduction and alkylation, the protein was purified in RP-HPLC as described below:

Column: Aquapore RP 300 (4.6×30 mm) cod. 0711-0055 Brownlee

Eluent A: 0.1 % aqueous TFA

Eluent B: 0.1 % TFA in CH₃CN

Column Temperature: + 40°C

5 UV detector set at 214 nm

Gradient				
Time (minutes)	Flow (ml/min)	% A	%B	Curve
0	1	95	5	
5	1	95	5	6
6	1	80	20	6
41	1	35	65	6
46	1	20	80	1
47	1	95	5	6
57	1	95	5	6

10

The purified material was dried in speed-vac, dissolved in 250 μ L of 0.1M ammonium bicarbonate pH 9,0 \pm 0.05 and incubated with 5 μ L of modified bovine trypsin at 37 \pm 1°C for 4 hours with intermittent shaking. The reaction was stopped by adding 60 μ L of 5% aqueous TFA.

15

ANALYTICAL RP-HPLC OF TRYPTIC PEPTIDE MAPPING

Half volume of the r-hIL-18BP peptide mixture was purified in RP-HPLC as described below:

Column: Waters Symmetry C18 3,5 μ m (4.6×75 mm)

20

Eluent A: 0.1 % aqueous TFA

Eluent B: 0.1 % TFA in CH₃CN

Temperature: +45°C

UV detector set at 214 nm

GRADIENT				
Time (minutes)	Flow (ml/min)	%A	%B	Curve
0	1.0	98	2	
2	1.0	98	2	6
61	1.0	57	43	6
63	1.0	10	90	1
65	1.0	98	2	6

EDMAN SEQUENCING ANALYSIS

Automated Edman sequencing was carried out on a Procise protein sequencer,
 5 according to the manufacturer's instructions.

MALDI-TOF

MALDI-ToF spectra were carried out on a Voyager PE-Pro, according to
 manufacturer instructions.

10

LC-ES/MS OF TRYPSIN PEPTIDE MAPPING

The r-hIL-18BP was submitted to the peptide mapping procedure following the
 procedure mentioned above. After the digestion an aliquot of the peptide mixture of
 each sample was analysed as described below:

15 Column: Waters Symmetry C18 3,5 μ m (4.6 \times 75 mm)
 Eluent A: 0.1 % aqueous TFA
 Eluent B: 0.1 % TFA in CH₃CN
 Temperature: +45°C
 UV detector set at 214 nm

20

Gradient				
Time (minutes)	Flow (ml/min)	%A	%B	Curve
0	0.7	98	2	

12	0.7	98	2	6
71	0.7	57	43	6
73	0.7	10	90	1
75	0.7	98	2	6

After UV detector the flow was split in order to introduce in the spectrometer source at 50 μ L/min.

The mass spectrometer has been set with the following parameters:

- 5 Capillary voltage: 3.5 KV
 Cone voltage: 35 V
 HV lenses: 0.45 KV
 Source temperature: 80°C
 Resolution: 14 HM; 14 LM

10

SEC SELECTED METHOD FOR DIMERS/AGGREGATES CONTENT

The SE-HPLC analysis was carried out as reported below:

Eluent	0.1 M H ₃ PO ₄ , 0.3 M Na ₂ SO ₄ , pH 7.3 with NaOH, CAN 3%
Column type	TSK G2000 SW _{XL} 7.8x300 code 08540
Autosampler temperature	+ 4°C \pm 2°C
Column temperature	Room temperature
Detection wavelength	214 nm
Flow rate of mobile phase	0.5 mL/min
Analysis time	30 minutes
Delay for next injection	Not less than 5 minutes

SDS-PAGE AND SILVER STAINING

- 15 Two micrograms of r-hIL18BP were loaded onto the precast gel ExcelGel® SDS Homogeneous 12.5% (by Amersham Biosciences) in non reducing conditions and run under constant voltage (600 V) at 15°C. Molecular weight markers and the Interim Reference Materials ST1P01/r-hIL-18BP were also loaded onto the gel.

- 20 After the electrophoretic run the gel was stained with the Silver Staining Kit-Protein (PlusOne) as described in the instructions contained in the kit leaflet. Briefly,

the gel was fixed for 30 minutes in a solution composed of acetic acid and ethanol. After a washing step, the sensitising solution was added and removed after 30 minutes. The gel was washed again and then reacted with the silver solution for 20 minutes. After a washing cycle, the staining was developed in developing solution and subsequently stopped. The gel was then thoroughly washed in water and kept in preserving solution before final storage in Cellophane Sheets.

Gels were scanned and data elaborated using the Phoretix 1D full software.

SDS-PAGE AND WESTERN BLOTTING

Two hundred nanograms of r-hIL-18BP were loaded onto the precast gel ExcelGel® SDS Homogeneous 12.5% (by Amersham Biosciences) in non reducing conditions and run under constant voltage (600 V) at 15°C. Molecular weight markers and the Interim Reference Materials ST1P01/r-hIL-18BP were also loaded onto the gel.

After the electrophoretic run, proteins were transferred from the gel onto a nitro-cellulose membrane by passive contact for 60 minutes at room temperature and probed with 0.1 µg/mL of the monoclonal antibody to r-hIL-18BP clone 582.10 (IPL). The reaction was revealed by a chemiluminescent substrate (ECL kit from Amersham Biosciences) after reaction with 1:2000 diluted goat anti-mouse IgG HRP conjugate. The light emission was detected by 10 seconds or 1 minute of exposure to a sensitive autoradiography film.

Coomassie blue or silver staining methods were adopted to detect the MW markers.

After the immunodetection, the film was scanned and the molecular weight (MW) values of the bands were automatically derived from the MW calibration curve using the Phoretix 1-D Full software.

KG-1 CELLS IN VITRO BIOASSAY

The biological activity of samples was quantified by using an in vitro bioassay. This bioassay was based on the ability of the human acute myelogenous leukemia cell line KG-1 to produce IFN-γ in response to human IL-18 plus human TNF-α in a dose-dependent manner. The r-hIL-18BP specifically binds r-hIL-18 neutralizing its biological activity thereby suppressing the production of IFN-γ.

Briefly, KG-1 cells at 1×10^5 cells/well were added to a 96 well plate already containing different concentrations of r-hIL-18BP in the presence of a fixed concentration of r-hIL18 (40 ng/ml in the well) plus a fixed concentration of r-hTNF- α (10 ng/ml in the well). The concentration of each of these two substances combined together was able to give the sub-maximal induction of production of IFN- γ on KG-1 cells. After 24 hr at 37°C, 5% CO₂, the plate was put at -20°C in order to submit the treated cells to a freeze/thaw cycle before performing the immunoassay to determine the quantity of IFN- γ present in the cell supernatant. The cell supernatants was collected and human IFN- γ measured by means of a specific immunoassay (ELISA h-IFN- γ , Duo Set R&D Systems kit). The amount of IFN- γ produced by the treated cells (either with standard curve or IL-18BP sample) was calculated by interpolating the y values (O.D.) on the IFN- γ Standard curve, provided with the kit, fitted by a Sigmoidal dose-response (4PL) Log/Log transformed, thus obtaining the x values (IFN- γ concentrations) (GraphPad Prism).

The biological activity of IL-18BP sample was determined vs the reference preparation by testing the sample at two concentrations falling in the linear part of the reference dose-response curve. At least two independent experiments were carried out. In each independent assay, each concentration was tested in dependent duplicates in a plate.

The titer of IL-18BP sample for each concentration tested, was calculated by interpolating the averaged (two replicates) y values (O.D.) of the amount of IFN- γ produced on the linear part of the reference dose-response curve (Log/Log transformed) thus obtaining the x values (IL-18BP activity).

The value obtained from each concentration was averaged and the final activity of IL-18BP drug substance sample was given by the arithmetic mean of the potencies obtained from each of the independent assay performed.

The titer of the different IL-18BP drug substances was calculated versus the Interim Reference Material ST1P01/r-hIL-18BP.

Two independent experiments were carried out.

RESULTS

Background

The primary structure of full length r-hIL-18BP is shown in Fig. 1. The protein has a C-terminal heterogeneity, with molecules ending at residue 164 (full length) and residue 163 (C-1aa), the latter being the main form. Mass spectrometric analysis of tryptic peptides has further shown that the molecule is highly glycosylated, carrying both N- and O-linked oligosaccharides.

The molecule contains four potential N-glycosylation sites, at Asn 49, Asn 64, Asn 73 and Asn 117. Only three of the four sites have been found glycosylated, i.e. Asn 49, Asn 73 and Asn 117, whereas Asn 64 has been found glycosylated only in trace amounts.

The average molecular weight of the whole molecule as determined by SDS-PAGE and SE-HPLC is approximately 50kDa.

The amino acid composition may be taken from table 4.

TABLE 4 Amino acid composition

Amino acid	Three letter code	Single letter code	No	%
Alanine	Ala	A	13	7.9%
Arginine	Arg	R	7	4.3%
Asparagine	Asn	N	4	2.4%
Aspartic acid	Asp	D	2	1.2%
Cysteine	Cys	C	6	3.7%
Glutamine	Gln	Q	12	7.3%
Glutamic acid	Glu	E	9	5.5%
Glycine	Gly	G	9	5.5%
Histidine	His	H	4	2.4%
Isoleucine	Ile	I	2	1.2%
Leucine	Leu	L	19	11.6%
Lysine	Lys	K	3	1.8%
Methionine	Met	M	0	0.0%
Phenylalanine	Phe	F	5	3.0%
Proline	Pro	P	17	10.4%
Serine	Ser	S	18	11.0%

29

Threonine	Thr	T	15	9.1%
Tryptophan	Trp	W	4	2.4%
Tyrosine	Tyr	Y	1	0.6%
Valine	Val	V	14	8.5%

The routine QC tests of batches from serum-free production revealed that:


- There was a non-conform peptide mapping profile (one major additional peak already during purification of reduced and alkylated protein);
- 5 • An abnormal SE-HPLC profile was obtained;
- A double band was detected in SDS-PAGE
- A similar profile was obtained in RP-HPLC
- The specific activity versus homogeneous IL-18BP produced in serum-containing medium (the "reference standard") was comparable.

10

PEPTIDE MAPPING PROCEDURE

Since r-hIL-18BP is a highly glycosylated molecule, presenting a high heterogeneity in terms of glycosylation, the protein was submitted to Neuraminidase treatment in order to reduce oligosaccharide heterogeneity due to sialic acid. The
15 protein was then submitted to reduction, carboxymethylation and purification in order to render the trypsin cleavage sites well accessible to the enzyme.

The peptide procedure was carried out with the following steps:



A chromatographic profile different to the one of r-hIL-18BP produced in serum-containing medium, was already detected during the purification of the reduced and alkylated r-hIL-18BP batches from serum-free production (not shown).

Furthermore, the peptide mapping profile of a truncated form of r-hIL-18BP, compared to the current reference standard r-hIL-18BP, showed both an extra peak and a different relative intensity of glycosylated peptides (not shown).

N-TERMINAL ANALYSIS

The sequence analysis of the intact molecule showed different fragments corresponding to molecule starting from residues 1, 16, 31, and in lower amounts from residues 69, 70, 107 and 125. The N-terminal analysis is depicted in Fig. 1.

MALDI-TOF

The spectra obtained by MALDI-TOF showed an additional peak at lower molecular weight (not shown).

SDS-PAGE analysis (Fig. 2)

The r-hIL-18BP has a relative molecular weight of about 50 kDa as assigned by 12.5% SDS-PAGE. Serum-free produced r-hIL-18BP showed an additional band of

about 40 kDa detected by silver staining. Both bands reacted with an IL-18BP-specific antibody (clone 582.10) in Western Blotting analysis. The silver stained SDS-PAGE gel is depicted in Fig. 2 A, the Western Blot in Fig. 2 B.

5 The lanes were occupied as follows:

- | | |
|---------------------------|-------------------------------|
| 6. MW marker | 5. MW marker |
| 7. r-hIL-18BP CT20 (2 µg) | 6. r-hIL-18BP CT20 (200 ng) |
| 8. r-hIL-18BP CT20 (2 µg) | 7. ST1P01/r-hIL-18BP (200 ng) |
| 9. r-hIL-18BP CT20 (2 µg) | 8. MW marker |
| 10. ST1P01/r-hIL-18BP | |

CT20 is a batch of truncated IL-18BP, while ST1P01 is the standard full-length IL-18BP without truncated forms.

10 BIOASSAY

It was assessed whether The results of specific activity of the different IL-18 BP drug substance batches are reported in table 5 where the untruncated (ILNCT16-18 and ST1P01) and truncated form (highlighted, ILNCT 19-22) are shown.

15

TABLE 5

IL-18BP bulks	Biological activity U/mL	Protein content by O.D. mg/mL	Specific activity U/mg
ILNCT16	1,005,906	60.3	16,682
ILNCT17	1,167,546	56.8	20,555
ILNCT18	1,150,841	55.3	20,811
ILNCT19	949,440	61.6	15,413
ILNCT20	1,225,693	57.2	21,428
ILNCT21	1,278,583	62.8	20,360

ILNCT22	1,347,902	59.8	22,540
ILNCT23	1,200,463	60.7	19,777
ILNCT24	1,013,834	56.65	17,896
ST1PO1	895,69	46.39 (AAA)	19,312

This experiment shows that truncated IL-18BP has a biological activity comparable to untruncated IL-18BP-

5 TRUNCATED R-HIL-18BP

In order to characterize the extra peak detected by different techniques, the SE-HPLC analysis was employed to separate the peaks of interest so as to submit them to further characterization steps. For the intended purpose the two peaks were collected separately.

10 In order to unequivocally identify the collected peaks, peak 1 and peak 2 were re-injected onto the HPLC column.

The two peaks were submitted to peptide mapping according to the protocol described above.

15 The chromatographic profiles of reduced and alkylated samples are reported in Fig. 3.

Only the main peaks of each fraction were submitted to peptide mapping and analysed by LC-ES/MS:

20 The extra peak (peptide 31-61) appearing in the peptide mapping profiles is due to internal cleavages of the molecule, as confirmed by the sequence analysis of the peaks and of the intact molecule.

Moreover the different intensities of glycosylated peptides (Pep. 1-15, Pep. 1-32 and Pep. 16-32) show a different glycosylation pattern.

The N-terminal analysis carried out onto peak 1 and peak 2 collected directly from

25 SE-HPLC analysis, gave the following results:

N-terminal analysis of Peak 1 isolated by SE-HPLC

N-term	T	P	V	S	Q	X	X	roughly 54%
From 31	A	K	Q	X	P	A	L	roughly 46%
From 16	S	T	K	D	P	C	P	trace

N-terminal analysis of Peak 2 isolated by SE-HPLC

From 16	S	T	K	D	P	C	P	roughly 63%
From 31	A	K	Q	X	P	A	L	roughly 37%
N-term	T	P	V	S	Q	X	X	trace

5

The apparent molecular weight assigned by SDS-PAGE (Fig. 3) was confirmed by MALDI-TOF spectra (not shown).

10

CONCLUSIONS

The results obtained employing different analytical tools showed that the following major cleavage sites can be identified:

- Protein truncated at residue 15, i.e. the sequence starting from residue 16 and ending at residue 163/164;
- 15 • Protein cleaved at residue 30, i.e. the full length sequence, from residue 1 to residue 163/164, with an internal clipping between residues 30 and 31, held together by disulfides;
- Protein both truncated at residue 15 and cleaved at residue 30, i.e. the sequence starting from residue 16 and ending at residue 163/164, with an internal clipping
- 20 between residues 30 and 31, held together by disulfides;

When a truncated form of r-hIL18BP is present, the above results show the following:

- A double band is detected by SDS-PAGE of samples. The two bands are detected
- 25 both by Silver staining and western blotting.
- The SE-HPLC analysis shows an anomalous profile.

- The RP-HPLC chromatographic profiles of reduced and alkylated samples is different as compared to the one of intact samples.
 - The peptide mapping profiles showed an extra peak.
 - The N-terminal sequence analysis confirms the presence of truncated forms of the molecule.
- 5
- Despite the truncated form is present, the specific activity is comparable to that of the intact r-hIL-18BP.

REFERENCES

1. Conti, B., J. W. Jahng, C. Tinti, J. H. Son, and T. H. Joh. 1997. Induction of interferon-gamma inducing factor in the adrenal cortex. *J. Biol. Chem.* 272:2035-2037.
- 5 2. DiDonato, J A, Hayakawa, M, Rothwarf, D M, Zandi, E and Karin, M. (1997), *Nature* 388, 16514-16517.
3. Engelmann, H., D. Novick, and D. Wallach. 1990. Two tumor necrosis factor-binding proteins purified from human urine. Evidence for immunological cross-reactivity with cell surface tumor necrosis factor receptors. *J. Biol. Chem.* 265:1531-1536.
- 10 4. Gracie JA, Robertson SE, McInnes IB *J Leukoc Biol* 2003 Feb;73(2):213-24
5. Kim SH, Eisenstein M, Reznikov L, Fantuzzi G, Novick D, Rubinstein M, Dinarello CA. Structural requirements of six naturally occurring isoforms of the IL-18 binding protein to inhibit IL-18. *Proc Natl Acad Sci U S A* 2000;97:1190-1195.
- 15 6. Micallef, M. J., T. Ohtsuki, K. Kohno, F. Tanabe, S. Ushio, M. Namba, T. Tanimoto, K. Torigoe, M. Fujii, M. Ikeda, S. Fukuda, and M. Kurimoto. 1996. Interferon-gamma-inducing factor enhances T helper 1 cytokine production by stimulated human T cells: synergism with interleukin-12 for interferon-gamma production. *Eur-J-Immunol* 26:1647-51 issn: 0014-2980.
- 20 7. Nakamura K, Okamura H, Wada M, Nagata K, Tamura T. *Infect Immun* 1989 Feb;57(2):590-5
8. Novick, D, Kim, S-H, Fantuzzi, G, Reznikov, L, Dinarello, C, and Rubinstein, M (1999). *Immunity* 10, 127-136.
9. Okamura H, Nagata K, Komatsu T, Tanimoto T, Nukata Y, Tanabe F, Akita K, Torigoe K, Okura T, Fukuda S, et al. *Infect Immun* 1995 Oct;63(10):3966-72
- 25 10. Rothe H, Jenkins NA, Copeland NG, Kolb H. *J Clin Invest* 1997 Feb 1;99(3):469-74
11. Yoshimoto T, Takeda, K, Tanaka, T, Ohkusu, K, Kashiwamura, S, Okamura, H, Akira, S and Nakanishi, K (1998), *J. Immunol.* 161, 3400-3407.
12. Xiang and Moss, *J. Biol. Chem.* 2001 276: 17380-6
- 30 13. Xiang and Moss, *J. Virol.* 2001 75 (20), 9947-54

CLAIMS

1. IL-18 binding protein (IL-18BP) comprising an amino acid sequence selected from
SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, or
5 SEQ ID NO: 7, but not comprising SEQ ID NO: 1, or a functional derivative, fusion
protein or salt thereof.
2. The IL-18 binding protein (IL-18BP) according to claim 1, less the C-terminal glycine
residue.
10
3. IL-18 binding protein (IL-18BP) consisting of the amino acid sequence of SEQ ID
NO: 1 and being N-glycosylated at Asn 49, Asn 73 and Asn 117, but not at Asn 64,
a functional derivative, fusion protein or salt thereof.
- 15 4. The IL-18 binding protein (IL-18BP) according to claim 1, less the C-terminal glycine
residue.
5. IL-18BP comprising a first polypeptide consisting of amino acids 1 to 30 of SEQ ID
NO: 1 and a second polypeptide consisting of amino acids 31 to 164 or of amino
20 acids 31 to 163, wherein the first and second polypeptide are linked by a disulfide
bond.
6. IL-18BP comprising a first polypeptide consisting of amino acids 15 to 30 of SEQ ID
NO: 1 and a second polypeptide consisting of amino acids 31 to 164 or of amino
25 acids 31 to 163, wherein the first and second polypeptide are linked by a disulfide
bond.
7. The IL-18BP according to any of the preceding claims, wherein the fused protein
comprises an immunoglobulin fusion.
30
8. The IL-18BP according to any of the preceding claims, wherein the functional
derivative comprises at least one moiety attached to one or more functional groups,
which occur as one or more side chains on the amino acid residues.

9. The IL-18BP according to claim 8, wherein the moiety is a polyethylene glycol (PEG) moiety.
- 5 10. A nucleic acid coding for an IL-18BP according to any of claims 1 to 7.
11. Host cell comprising a nucleic acid according to claim 10.
- 10 12. Process for production of an IL-18BP according to any of claims 1 to 7 comprising the step of culturing a host cell according to claim 11 under conditions suitable for expression of said IL-18BP.
- 15 13. Process for production of an IL-18BP according to any of claims 1 to 7 comprising the step of isolating the IL-18BP from the cell culture supernatant of a host cell according to claim 11.
14. Composition comprising an IL-18BP according to any of claims 1 to 9.
- 20 15. Use of an IL-18BP according to any of claims 1 to 9 for the preparation of a medicament for treatment and/or prevention of an IL-18 mediated disease.
- 25 16. Use according to claim 15, wherein the IL-18 mediated disease is selected from: psoriasis, arthritis, in particular rheumatoid arthritis, inflammatory bowel disease, in particular Crohn's disease, liver injury, atherosclerosis, sepsis, myocardial infarction, traumatic brain injury, allergy, peripheral vascular disease, multiple sclerosis.
- 30 17. The use according to claim 15 or 16, wherein the medicament further comprises an interferon, for simultaneous, sequential or separate use.
18. The use according to claim 17, wherein the interferon is interferon- β .

19. The use according to claim 15 or 16, wherein the medicament further comprises an inhibitor of Tumor Necrosis Factor (TNF) for simultaneous, sequential or separate use.
- 5 20. The use according to claim 19, wherein the inhibitor of TNF is a soluble TNF receptor.
21. The use according to claims 15 to 20, wherein IL-18BP is used in an amount of about 0.001 to 1000 mg/kg of body weight, or about 0.01 to 100 mg/kg of body weight or about 0.1 to 10 mg/kg of body weight or about 5 mg/kg of body weight.
- 10 22. The use according to any of claims 15 to 21, wherein the IL-18BP variant is administered subcutaneously.
- 15 23. The use according to any of claims 15 to 21, wherein the IL-18BP variant is administered intramuscularly.
24. Use of an expression vector comprising a nucleic acid according to claim 9 for the manufacture of a medicament for the treatment and/or prevention of an IL-18 mediated disease.
- 20 25. Use of a cell that has been genetically modified to produce an IL-18BP according to any of claims claims 1 to 7 for the manufacture of a medicament for the treatment and/or prevention of an IL-18 mediated disease.




FIG. 1





Fig. 2



SEQUENCE LISTING

5 <110> Applied Research Systems ARS Holding N.V:

10 <120> Active fragments of the IL-18 binding protein

<130> 818 WO

15

<160> 7

20

<170> PatentIn version 3.1

25 <210> 1

<211> 164

<212> PRT

30

<213> homo sapiens

35 <400> 1

Thr Pro Val Ser Gln Thr Thr Thr Ala Ala Thr Ala Ser Val Arg Ser
1 5 10 15

40

Thr Lys Asp Pro Cys Pro Ser Gln Pro Pro Val Phe Pro Ala Ala Lys
20 25 30

45 Gln Cys Pro Ala Leu Glu Val Thr Trp Pro Glu Val Glu Val Pro Leu
35 40 45

50 Asn Gly Thr Leu Ser Leu Ser Cys Val Ala Cys Ser Arg Phe Pro Asn
50 55 60

Phe Ser Ile Leu Tyr Trp Leu Gly Asn Gly Ser Phe Ile Glu His Leu
 65 70 75 80

5 Pro Gly Arg Leu Trp Glu Gly Ser Thr Ser Arg Glu Arg Gly Ser Thr
 85 90 95

10 Gly Thr Gln Leu Cys Lys Ala Leu Val Leu Glu Gln Leu Thr Pro Ala
 100 105 110

15 Leu His Ser Thr Asn Phe Ser Cys Val Leu Val Asp Pro Glu Gln Val
 115 120 125

20 Val Gln Arg His Val Val Leu Ala Gln Leu Trp Ala Gly Leu Arg Ala
 130 135 140

Thr Leu Pro Pro Thr Gln Glu Ala Leu Pro Ser Ser His Ser Ser Pro
 145 150 155 160

25 Gln Gln Gln Gly

30 <210> 2
 <211> 149
 <212> PRT

35 <213> homo sapiens

40 <400> 2
 Ser Thr Lys Asp Pro Cys Pro Ser Gln Pro Pro Val Phe Pro Ala Ala
 1 5 10 15

45 Lys Gln Cys Pro Ala Leu Glu Val Thr Trp Pro Glu Val Glu Val Pro
 20 25 30

50 Leu Asn Gly Thr Leu Ser Leu Ser Cys Val Ala Cys Ser Arg Phe Pro
 35 40 45

5 Asn Phe Ser Ile Leu Tyr Trp Leu Gly Asn Gly Ser Phe Ile Glu His
 50 55 60

10 Leu Pro Gly Arg Leu Trp Glu Gly Ser Thr Ser Arg Glu Arg Gly Ser
 65 70 75 80

15 Thr Gly Thr Gln Leu Cys Lys Ala Leu Val Leu Glu Gln Leu Thr Pro
 85 90 95

20 Ala Leu His Ser Thr Asn Phe Ser Cys Val Leu Val Asp Pro Glu Gln
 100 105 110

25 Val Val Gln Arg His Val Val Leu Ala Gln Leu Trp Ala Gly Leu Arg
 115 120 125

30 Ala Thr Leu Pro Pro Thr Gln Glu Ala Leu Pro Ser Ser His Ser Ser
 130 135 140

35 Pro Gln Gln Gln Gly
 145

40 <210> 3
 <211> 134
 <212> PRT
 <213> homo sapiens

45 Ala Lys Gln Cys Pro Ala Leu Glu Val Thr Trp Pro Glu Val Glu Val
 1 5 10 15

50 Pro Leu Asn Gly Thr Leu Ser Leu Ser Cys Val Ala Cys Ser Arg Phe
 20 25 30

55 Pro Asn Phe Ser Ile Leu Tyr Trp Leu Gly Asn Gly Ser Phe Ile Glu

35 40 45
 5 His Leu Pro Gly Arg Leu Trp Glu Gly Ser Thr Ser Arg Glu Arg Gly
 50 55 60
 10 Ser Thr Gly Thr Gln Leu Cys Lys Ala Leu Val Leu Glu Gln Leu Thr
 65 70 75 80
 15 Pro Ala Leu His Ser Thr Asn Phe Ser Cys Val Leu Val Asp Pro Glu
 85 90 95
 20 Gln Val Val Gln Arg His Val Val Leu Ala Gln Leu Trp Ala Gly Leu
 100 105 110
 25 Arg Ala Thr Leu Pro Pro Thr Gln Glu Ala Leu Pro Ser Ser His Ser
 115 120 125
 30 Ser Pro Gln Gln Gln Gly
 130
 35 <210> 4
 40 <211> 96
 <212> PRT
 <213> homo sapiens
 45 <400> 4
 50 Tyr Trp Leu Gly Asn Gly Ser Phe Ile Glu His Leu Pro Gly Arg Leu
 1 5 10 15
 Trp Glu Gly Ser Thr Ser Arg Glu Arg Gly Ser Thr Gly Thr Gln Leu
 20 25 30
 Cys Lys Ala Leu Val Leu Glu Gln Leu Thr Pro Ala Leu His Ser Thr
 35 40 45

Asn Phe Ser Cys Val Leu Val Asp Pro Glu Gln Val Val Gln Arg His
50 55 60

5 Val Val Leu Ala Gln Leu Trp Ala Gly Leu Arg Ala Thr Leu Pro Pro
65 70 75 80

10 Thr Gln Glu Ala Leu Pro Ser Ser His Ser Ser Pro Gln Gln Gln Gly
85 90 95

<210> 5

15 <211> 95

<212> PRT

<213> homo sapiens

20

<400> 5

25 Trp Leu Gly Asn Gly Ser Phe Ile Glu His Leu Pro Gly Arg Leu Trp
1 5 10 15

30 Glu Gly Ser Thr Ser Arg Glu Arg Gly Ser Thr Gly Thr Gln Leu Cys
20 25 30

35 Lys Ala Leu Val Leu Glu Gln Leu Thr Pro Ala Leu His Ser Thr Asn
35 40 45

Phe Ser Cys Val Leu Val Asp Pro Glu Gln Val Val Gln Arg His Val
50 55 60

40 Val Leu Ala Gln Leu Trp Ala Gly Leu Arg Ala Thr Leu Pro Pro Thr
65 70 75 80

45 Gln Glu Ala Leu Pro Ser Ser His Ser Ser Pro Gln Gln Gln Gly
85 90 95

<210> 6

50

<211> 58

<212> PRT

5 <213> homo sapiens

<400> 6

10 Glu Gln Leu Thr Pro Ala Leu His Ser Thr Asn Phe Ser Cys Val Leu
1 5 10 15

15 Val Asp Pro Glu Gln Val Val Gln Arg His Val Val Leu Ala Gln Leu
20 25 30

20 Trp Ala Gly Leu Arg Ala Thr Leu Pro Pro Thr Gln Glu Ala Leu Pro
35 40 45

25 Ser Ser His Ser Ser Pro Gln Gln Gln Gly
50 55

<210> 7

<211> 40

30 <212> PRT

<213> homo sapiens

35

<400> 7

40 Pro Glu Gln Val Val Gln Arg His Val Val Leu Ala Gln Leu Trp Ala
1 5 10 15

45 Gly Leu Arg Ala Thr Leu Pro Pro Thr Gln Glu Ala Leu Pro Ser Ser
20 25 30

His Ser Ser Pro Gln Gln Gln Gly
35 40

50

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP2004/005073

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07K14/715 C07K14/54 C07K14/47 A61K38/20

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07K A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, MEDLINE, CHEM ABS Data, EMBASE, BIOSIS, EMBL

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 99/09063 A (RUBINSTEIN MENACHEM ;KIM SOO HYUN (IL); NOVICK DANIELA (IL); YEDA) 25 February 1999 (1999-02-25) cited in the application the whole document	3,7-16
X	EP 1 110 969 A (HAYASHIBARA BIOCHEM LAB) 27 June 2001 (2001-06-27)	1,10-16, 19,22,23
Y	SEQ. ID. NO: 36 contains SEQ. ID. NO: 7 the whole document	20,24,25
	----- -/-	

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

17 September 2004

Date of mailing of the international search report

29/09/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Armandola, E

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP2004/005073

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	XIANG Y ET AL: "Identification of Human and Mouse Homologs of the MC51L-53L-54L Family of Secreted Glycoproteins Encoded by the Molluscum Contagiosum Poxvirus" VIROLOGY, ACADEMIC PRESS, ORLANDO, US, vol. 257, no. 2, 10 May 1999 (1999-05-10), pages 297-302, XP004439941 ISSN: 0042-6822 the whole document	10-14
X	YASUSHI A ET AL: "CLONING AND EXPRESSION OF INTERLEUKIN-18 BINDING PROTEIN" FEBS LETTERS, ELSEVIER SCIENCE PUBLISHERS, AMSTERDAM, NL, vol. 445, 1999, pages 338-342, XP002180254 ISSN: 0014-5793 the whole document	3, 8, 10-14
X	WO 02/32374 A (BORN TERESA L ; SIMS JOHN E (US); IMMUNEX CORP (US); MOHLER KENDALL) 25 April 2002 (2002-04-25)	7, 10-16, 21
Y	the whole document	20, 24, 25
X	NOVICK D ET AL: "INTERLEUKIN-18 BINDING PROTEIN: A NOVEL MODULATOR OF THE TH1 CYTOKINE RESPONSE" IMMUNITY, CELL PRESS, US, vol. 10, no. 1, January 1999 (1999-01), pages 127-136, XP000973488 ISSN: 1074-7613 cited in the application	3
Y	the whole document	10, 11
X	KIM S-H ET AL: "STRUCTURAL REQUIREMENTS OF SIX NATURALLY OCCURRING ISOFORMS OF THE IL-18 BINDING PROTEIN TO INHIBIT IL-18" PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA, NATIONAL ACADEMY OF SCIENCE. WASHINGTON, US, vol. 97, no. 3, 1 February 2000 (2000-02-01), pages 1190-1195, XP000941910 ISSN: 0027-8424 cited in the application	3
Y	the whole document	10, 11

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP2004/005073

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9909063	A	25-02-1999	AU 755794 B2	19-12-2002
			AU 8746098 A	08-03-1999
			BG 104149 A	31-08-2000
			BR 9811940 A	05-09-2000
			CA 2298855 A1	25-02-1999
			CN 1267307 T	20-09-2000
			DE 1003781 T1	11-01-2001
			EA 3675 B1	28-08-2003
			EE 200000073 A	16-10-2000
			EP 1003781 A1	31-05-2000
			ES 2149149 T1	01-11-2000
			HU 0003533 A2	29-01-2001
			WO 9909063 A1	25-02-1999
			JP 2001514876 T	18-09-2001
			NO 20000700 A	14-04-2000
			NZ 502392 A	20-12-2002
			PL 338647 A1	06-11-2000
			SK 1762000 A3	12-09-2000
			US 2004037829 A1	26-02-2004
			US 6605280 B1	12-08-2003
			ZA 9807310 A	14-08-2000
EP 1110969	A	27-06-2001	BR 9816013 A	21-01-2003
			CA 2340579 A1	09-03-2000
			EP 1110969 A1	27-06-2001
			WO 0012555 A1	09-03-2000
			JP 2000210087 A	02-08-2000
WO 0232374	A	25-04-2002	AU 2441702 A	29-04-2002
			WO 0232374 A2	25-04-2002
			US 2002098185 A1	25-07-2002